



National Neonatal Audit Programme

A guide to the 2020 audit measures

November 2019, v1.0

National Neonatal Audit Programme (NNAP): A guide to the 2020 audit measures

This document sets out the details of the 2020 NNAP audit measures and describes the following details for each measure:

- NNAP audit measure
- Changes to the audit measure for 2020 data
- NNAP standard and source of standard
- Inclusion criteria
- Attributing results
- Deriving outcomes
- Where the required data should be entered on BadgerNet

Audit measure	Page
About the NNAP dataset and methodology	4
Antenatal steroids Is a mother who delivers a baby between 23 and 33 weeks gestational age inclusive given at least one dose of antenatal steroids?	5
Antenatal magnesium sulphate Is a mother who delivers a baby below 30 weeks gestational age given magnesium sulphate in the 24 hours prior to delivery?	7
Birth in a centre with a neonatal intensive care unit (NICU) Is an admitted baby born at less than 27 weeks gestational age delivered in a maternity service on the same site as a designated NICU?	9
Deferred cord clamping Does a baby born at less than 32 weeks gestational age have their cord clamped at or after one minute?	11
Promoting normal temperature on admission for very preterm babies Does an admitted baby born at less than 32 weeks gestational age have a first temperature on admission which is both between 36.5–37.5°C and measured within one hour of birth?	11
Parental consultation within 24 hours of admission Is there a documented consultation with parents by a senior member of the neonatal team within 24 hours of admission?	15
Parental presence at consultant ward rounds For a baby admitted for more than 24 hours, did at least one parent attend a consultant ward round?	17
On-time screening for retinopathy of prematurity Does an admitted baby born weighing less than 1501 g, or at gestational age of less than 32 weeks, undergo the first retinopathy of prematurity (ROP) screening in accordance with the NNAP interpretation of the current guideline recommendations?	19
Bloodstream infection Does an admitted baby have one or more episodes of bloodstream infection, characterised by one or more positive blood cultures taken, after 72 hours of age?	22

<p>Central line associated bloodstream infection (QISD)</p> <p>How many babies have a positive blood culture (any species) with a central line present, after the first 72 hours of life, per 1000 central line days?</p>	24
<p>Bronchopulmonary dysplasia (BPD)</p> <p>Does an admitted baby born at less than 32 weeks develop bronchopulmonary dysplasia (BPD)?</p>	26
<p>Necrotising enterocolitis</p> <p>Does an admitted baby born at less than 32 weeks gestational age meet the NNAP surveillance definition for necrotising enterocolitis (NEC) on one or more occasion?</p>	28
<p>Minimising inappropriate separation of mother and term baby</p> <p>For a baby born at greater than or equal to 37 weeks gestational age, who did not have any surgery or a transfer during any admission, how many special care^a or normal care^b days were provided when oxygen was not administered?</p> <p><i>(a= Healthcare Resource Group (HRG) 3 or b= HRG 5, as defined by the NHS England neonatal critical care service specification)</i></p>	31
<p>Minimising inappropriate separation of mother and late preterm baby</p> <p>For a baby born at 34-36 weeks gestational age, who did not have any surgery or a transfer during any admission, how many special care ^a or normal care^b days were provided when oxygen was not administered?</p> <p><i>(a= HRG 3 or b= HRG 5, as defined by the NHS England neonatal critical care service specification)</i></p>	34
<p>Early breastmilk feeding</p> <p>Does a baby born at less than 32 weeks gestational age receive any of their own mother's milk at day 14 of life?</p>	37
<p>Breastmilk feeding at discharge home</p> <p>Does a baby born at less than 32 weeks gestational age receive any of their own mother's milk at discharge to home from a neonatal unit?</p>	39
<p>Follow-up at two years of age</p> <p>Does a baby born at less than 30 weeks gestational age receive medical follow-up at two years gestationally corrected age (18-30 gestationally corrected age range of acceptable ages)?</p> <p>Does a baby have complete results of a structured assessment recorded?</p>	41
<p>Mortality to discharge in very preterm babies</p> <p>Does a baby born at less than 32 weeks gestational age die before discharge home, or 44 weeks post-menstrual age (whichever occurs sooner)?</p>	46
<p>Nurse staffing on neonatal units</p> <p>Measure one: What proportion of nursing shifts are numerically staffed according to guidelines and service specification?</p> <p>Measure two: What proportion of shifts have sufficient staff qualified in speciality (QIS)?</p> <p>Measure three: How many additional nursing shifts are required to be worked to meet guidelines and service specification?</p>	49

About the NNAP dataset and methodology

Data for the NNAP analyses are extracted from the National Neonatal Research Database (NNRD) held at the Neonatal Data Analysis Unit (NDAU). The NNRD contains a predefined set of variables (the National Neonatal Dataset) obtained from the electronic neonatal patient records of each participating NHS trust or health board. Data are downloaded from the BadgerNet patient record system used in neonatal units and transferred to NDAU with health board and trust Caldicott Guardian approval. In Scotland, a separate approval was received from the Public Benefit and Privacy Panel for health and social care.

Inclusion criteria

The following inclusion criteria apply to all NNAP measures:

- Babies who were admitted for neonatal care
- Babies who had care provided by an NNAP unit
- Babies whose parents or carers have not opted them out of secondary use of their data

Further criteria specific to each audit measure can be found in the measure descriptions.

Case ascertainment

In usual practice, every baby admitted to a participating neonatal unit is entered on the BadgerNet patient record system, and is eligible for inclusion in the NNAP; the audit therefore achieves 100% case ascertainment in the participating organisations. Babies receiving special care in transitional care areas or postnatal wards can also be entered, but it is known that some units do not enter data for such babies and for this reason measures do not concentrate on care outside neonatal units.

Data collection period

Unless otherwise stated, the cohort of babies included in the 2020 audit year are babies who experience their final neonatal discharge in the calendar year 1 January 2020 to 31 December 2020. There are some exceptions to this; the minimising separation of mother and baby (term and late preterm) measures use birth year, follow-up at two years of age comprises babies born between July 2017 and June 2018, and mortality until discharge home comprises babies born between July 2017 and June 2020.

Describing gestational age

Gestation refers to “completed weeks” of gestation, based on the best available estimate recorded by clinical staff in BadgerNet. Gestation is typically described in bands – where we refer to all babies born at less than 32 weeks we mean all babies with a gestation at birth of up to 31 weeks and six days inclusive. Where we refer to 23-33 weeks inclusive, we mean to include all babies with a gestation at birth of 23 weeks and 0 days to babies with a gestation at birth of 33 weeks and 6 days.

Antenatal steroids

Is a mother who delivers a baby between 23 and 33 weeks gestational age inclusive given at least one dose of antenatal steroids?

Change to audit measure for the 2020 data year: The NNAP does not intend to actively report this measure in the future, however we include the measure here to encourage continued data entry and quality improvement activity around this measure. Please continue to monitor your performance and data entry for this measure.

NNAP standard

Developmental standard: Eighty-five percent (85%) of mothers should receive at least one dose of antenatal steroids.

Source of standard: NNAP Project Board

Comparison standard for outlier analysis: National rate

Inclusion criteria

- Babies who experienced their final neonatal discharge in the calendar year of analysis.
- Babies with gestational age at birth between 23 and 33 weeks inclusive.
- Only data from the first known episode of care will be considered for analysis.
- For multiple births, only one baby will be included so that each mother is only counted once per delivery.

Attribution

Results will be reported for each hospital of birth and network of birth. When the place of birth is unknown, a non-hospital location, or a hospital without a neonatal unit, the place of birth will be assigned as *Other*.

Deriving outcomes

NNAP will use data from steroids given and steroid courses to determine if the baby’s mother received any antenatal steroids. When these two fields contain contradictory information then “the most positive”* data will be assumed correct (Table 1). When multiple births present different data, data will be taken from the baby with “the most positive” outcome will be considered for the audit.

Table 1: Categorising data for antenatal steroids from BadgerNet fields Steroids given and Steroid courses

Steroids given	Steroid courses		
	Complete course	Incomplete course	Unknown / missing data
Yes	Steroids given	Steroids given	Steroids given
No	Steroids given	Steroids given	Steroids not given
Unknown / missing data	Steroids given	Steroids not given	Missing / unknown data

*Most to least positive: steroids given, steroids not given, unknown / missing data.

BadgerNet data source

Data from the *Pregnancy Details* page, for *Steroids during pregnancy* will be used to determine if antenatal steroids were given.

The screenshot displays the 'Pregnancy Details' page in BadgerNet, with a sidebar on the left containing navigation links: 'Pregnancy Details', 'Pregnancy To Unit', 'Parent Details', 'Pregnancy Details', 'Labour and Delivery', 'P and Professionals', 'RBS II', and 'Final Neonatal Outcome'. The main content area is divided into three sections:

- Antenatal screening:** Contains radio button options for Rubella Status (Immune, Non-immune, Not tested, Unknown), Hepatitis B surface antigen status (Negative, Positive, Not tested, Unknown), HIV Status (Negative, Positive, Not tested, Unknown), Syphilis (Negative, Positive, Not tested, Unknown), VDRL (Negative, Positive, Not tested, Unknown), TPHA (Negative, Positive, Not tested, Unknown), Hepatitis C virus antibody (Negative, Positive, Not tested, Unknown), and Hepatitis C virus PCR (Negative, Positive, Not tested, Unknown). It also includes three dropdown menus for 'Other screening 1', 'Other screening 2', and 'Other screening 3'.
- Antenatal:** Includes a radio button for 'Received Antenatal Care' (No, Yes, Unknown). Under the 'Dates' sub-section, there are dropdown menus for 'Date of First Ultrasound scan', 'Last menstrual period', 'EDD from LMP', and 'Agreed EDD', along with radio buttons for 'Calculated gestation' (Weeks, Days). Below this are dropdown menus for 'Detailed Anomaly Scan' and 'Doppler studies', each followed by a text area for 'Comments'.
- Steroids during pregnancy:** Features radio buttons for 'Steroids given' (Yes, No, Unknown), a dropdown for 'Last dose' (at), a dropdown for 'Courses given', and radio buttons for 'Which' (Betamethasone, Dexamethasone). Two red arrows point to the 'Steroids given' and 'Courses given' fields.

Figure 1: Pregnancy Details page, BadgerNet

Antenatal magnesium sulphate

Is a mother who delivers a baby below 30 weeks gestational age given magnesium sulphate in the 24 hours prior to delivery?

Change to audit measure for 2020 data year: None.

NNAP standard

Developmental standard: Eighty-five percent (85%) of eligible mothers should receive antenatal magnesium sulphate.

Source of standard: PReCePT¹.

Comparison standard for outlier analysis: National rate

Inclusion criteria

- Babies who experienced their final neonatal discharge in the calendar year of analysis.
- Babies with gestational age at birth less than 30 weeks.
- Only the first known episode of care will be considered for analysis.
- For multiple births, only one baby will be included so that each mother is only counted once per delivery.

Attribution

Results will be reported for each hospital of birth and network of birth. When the place of birth is unknown, a non-hospital location, or a hospital without a neonatal unit, the place of birth will be assigned as "Other".

Deriving outcomes

NNAP will use data from *Mother received magnesium sulphate in 24 hours prior to delivery* to categorise the administration of magnesium sulphate. When multiple births present different data, data will be taken from the baby with "the most positive"* for the audit (Table 2).

Table 2: Categorising antenatal magnesium sulphate from BadgerNet field *Mother received magnesium sulphate in 24 hours prior to delivery*

Mother received magnesium sulphate in 24 hours prior to delivery	NNAP category
Yes	Magnesium sulphate given
No	Magnesium sulphate not given
Missing / unknown	Missing / unknown data

*Most to least positive: magnesium sulphate given, magnesium sulphate not given, missing / unknown data.

BadgerNet data source

Data from the *Labour and Delivery* page for Mother received magnesium sulphate in 24 hours prior to delivery will be used to determine if magnesium sulphate was given.

The screenshot shows the 'Labour and Delivery' page in BadgerNet. The left sidebar contains navigation tabs: Baby Details, Admission To Unit, Parent Details, Pregnancy Details, Labour and Delivery (highlighted), GP and Professionals, CRIB II, and Final Neonatal Outcome. The main content area is divided into two sections: 'Labour' and 'Delivery'. In the 'Labour' section, the field 'Mother received Magnesium Sulphate in 24 hours prior to delivery' is highlighted in blue and has a red arrow pointing to it. The value for this field is 'No'. Other fields in the 'Labour' section include: Onset of labour (radio buttons for Spontaneous, Induced, None), Meconium Stained Liquor (radio buttons for Yes, No), Labour history (text input), Drugs in Labour (dropdown), Date and Time Membranes Ruptured (dropdown for date and time), Duration of Membrane Rupture (text input for hours), Maternal pyrexia in labour more than 38C (radio buttons for No, Yes, Unknown), Intrapartum antibiotics given (radio buttons for No, Yes), and Reason Magnesium Sulphate not given (radio buttons for Not offered, Not appropriate, Delivery imminent (within 1 hour), Contraindicated, Mother declined). The 'Delivery' section includes: Presentation Immediately Before Delivery (dropdown), Mode of Delivery (radio buttons for Emergency caesarean - not in labour, Emergency caesarean - in labour, Elective section - not in labour, Elective section - in labour, Vaginal - forceps assisted, Vaginal - spontaneous, Vaginal - ventouse assisted), Baby delivered in water (radio buttons for Yes, No), Condition at birth (text input), Offensive liquor? (radio buttons for No, Yes, Unknown), Apgar Score at 1 min, 5 mins, 10 mins, and 20 mins (radio buttons for scores 0-10), and Was cord clamping immediate (radio buttons for Yes, No, Unknown) with a time from birth to clamp field in minutes and seconds.

Figure 2: Labour and delivery page, BadgerNet

Additional information

Strong systematic review evidence of more than 6000 babies in meta-analysed randomised controlled trials suggests that antenatal magnesium sulphate therapy reduces the risk of cerebral palsy by about 30%, with benefits seen regardless of the reason for administration². Royal College of Obstetricians and Gynaecologists guidance supports its use³.

Birth in a centre with a neonatal intensive care unit (NICU)

Is an admitted baby born at less than 27 weeks gestational age delivered in a maternity service on the same site as a designated NICU⁴?

Change to audit measure for 2020 data year: None.

NNAP standard

Developmental standard: Eighty-five (85%) of babies born at less than 27 weeks gestational age should be delivered in a maternity service on the same site as a NICU.

Source of standard: Neonatal Critical Care Clinical Reference Group, NHS England.

Comparison standard for outlier analysis: National rate

Inclusion criteria

- Babies who experienced their final neonatal discharge in the calendar year of analysis.
- Babies with gestational age at birth of less than 27 weeks.
- Only data from the first known episode of care will be considered for analysis.
- For multiple births, only one baby will be included so that each mother is only counted once per delivery.

Attribution

Attribution will be to the neonatal network of birth. Outcomes will be reported by neonatal network of birth. When mothers deliver in a location not associated with a neonatal network (home, in transit, unknown, etc.) they will be assigned to a neonatal network based on the hospital of first admission.

Deriving outcomes

NNAP will use data from the place of birth to determine the level of neonatal unit within the hospital where babies were delivered (Table 3).

Table 3: Categorising birth in a centre with a NICU from BadgerNet field *Place of birth*

Place of birth	NNAP category
Hospital with a designated NICU	Delivered in appropriately designated location (NICU)
Hospital with a designated local neonatal unit (LNU) / special care unit (SCU)	Delivered in inappropriately designated location (LNU / SCU)
Hospital with no designated neonatal unit/ non-hospital location	Delivered in location without neonatal facilities

BadgerNet data source

Data on place of birth are captured on BadgerNet on the *Details at Birth and Admission* page. Values from the *Place of birth* field are used by NNAP for analysis (Figure 3).

The screenshot shows the 'Details at Birth and Admission' page in BadgerNet. On the left is a navigation menu with categories like 'Baby Details', 'Admission To Unit', 'Parent Details', 'Pregnancy Details', 'Labour and Delivery', 'GP and Professionals', 'CRIB ID', and 'Final Neonatal Outcome'. The main content area is split into two sections. The top section, 'Baby's Identification', contains fields for NHS Number (T:WDTJ2CFXGC), a 'Create temporary NHS Number' button, Baby's Local Hospital ID, Badger ID (AAZ229), Surname (BABYDJARYTEST3), Forename (BABYDJARYTEST3), Other/Previous Surnames, Sex (Female selected), and Non NHS patient status (Yes selected). The bottom section, 'General information', includes Birth Order (1 of 1), Date and Time of Birth (14 Apr 16 at 03:30), Place of birth (Test Hospital A), Code (XX888), a 'Place of birth is this hospital' button, Birth Location, and Agreed gestation at birth (Weeks and Days). A red arrow points to the 'Place of birth' field.

Figure 3: Details at birth and admission page, BadgerNet

Additional information

Background

Under the service specification⁴, neonatal networks intend to concentrate the delivery of babies to be born at less than 27 weeks gestation in units configured to deliver their care, i.e. a NICU. Evidence suggests that outcomes are improved by providing the care of the most vulnerable babies in units with a higher turnover, and minimising postnatal transfers.

Opportunities for quality improvement

Rates of delivery at less than 27 weeks appear to vary by neonatal network, and have changed over time. Ongoing "transformation" of specialist neonatal care may result in further modifications of patient pathways. In networks where a significant number of babies are delivered in centres without a NICU on site, there may be a quality improvement opportunity.

Deferred cord clamping for very preterm babies

Does a baby born at less than 32 weeks gestational age have their cord clamped at or after one minute?

Change to audit measure for 2020 data year: New measure.

NNAP standard

Developmental standard: Benchmarking only.

Comparison standard for outlier analysis: Not applicable, no outlier analysis.

Inclusion criteria

- Babies with gestational age at birth of less than 32 weeks.
- Babies who were admitted to a neonatal unit.
- Babies who experienced their final neonatal discharge in the calendar year of analysis.
- Only the first known episode of care will be considered for analysis.

Attribution

Results will be reported for each hospital of birth and network of birth. When the place of birth is unknown, a non-hospital location, or a hospital without a neonatal unit, the place of birth will be assigned as “Other”.

Deriving outcomes

NNAP will use data from the *Time of Cord Clamping* to determine whether cord clamping was deferred for 1 minute or more (Table 4).

Table 4: Categorising time of cord clamping in the NNAP

Time of cord clamping				
Missing	< 0 minutes	Less than a minute after birth	1 - 2 minutes after birth	More than 2 minutes after birth
Time of cord clamping missing	Time of cord clamping missing	Less than 1 minute after birth	Deferred cord clamping (≥ 1 minute)	Deferred cord clamping (≥ 1 minute)

BadgerNet data source

Data on time of cord clamping are taken from the field *Time from birth to clamp* on the *Labour and Delivery* page.

The screenshot shows the 'Delivery' section of a medical record form. The 'Presentation Immediately Before Delivery' is 'Cephalic - unspecified'. The 'Mode of Delivery' is 'Vaginal - spontaneous'. The 'Baby delivered in water' is 'No'. The 'Condition at birth' is 'Born in reasonable condition'. The 'Offensive liquor?' is 'Yes'. The 'Apgar Score at 1 min', 'Apgar Score at 5 mins', and 'Apgar Score at 10 mins' are all '10'. The 'Was cord clamping immediate' field is 'Yes', and the 'Time from birth to clamp' is '40' seconds. Two red arrows point to the 'Was cord clamping immediate' and 'Time from birth to clamp' fields.

Additional information

Background

The timing of clamping of the umbilical cord at birth was first considered time critical as part of a triad of measures to reduce maternal post-partum bleeding in the 1960s.⁵ However, it is now known that immediate cord clamping is not necessary for this reason, and more importantly has been shown to be harmful on aggregate in preterm infants.^{6,7} Recent trials, and an important meta-analysis including data from 2834 infants, show that avoiding immediate cord clamping reduces mortality by 32%.^{6,8} It is clear that within this evidence base, and therefore in ongoing clinical practice, that because deferred cord clamping was not practised in all infants randomised to this approach, it is unknown if changing to immediate cord clamping for a subgroup of infants who appear non vigorous at birth is appropriate. For example in one large trial, 73% of infants randomised to deferral received deferral.⁷ International clinical guidelines have recommended deferred cord clamping for some time.⁹ Cord milking is excluded from this measure - trial data suggest this is harmful.

Opportunities for Quality Improvement

Trials have shown a very large reduction in mortality. Deferred cord clamping has been shown to be incompletely implemented in the UK. NNAP audit users have suggested this is an important outcome that could be measured, and that measurement might facilitate adoption of deferral by benchmarking and review of practise. The focus on very preterm infants (those less than 32 weeks gestation) is on account of the higher mortality in this group, and to be confident that all infants delivered within a hospital are admitted to a neonatal unit, and are thus captured within the measure.

Promoting normal temperature on admission for very preterm babies

Does an admitted baby born at less than 32 weeks gestational age have a first temperature on admission which is both between 36.5–37.5°C and measured within one hour of birth^{4,10}?

Change to audit measure for 2020 data year: None.

NNAP standard

Developmental standard: First temperature on admission should be taken within an hour of birth for all eligible babies.

The composite measure of timeliness and normal temperature should be met for at least ninety percent (90%) of babies.

Source of standard: NNAP Project Board

Comparative standard for outlier analysis: National rate.

Inclusion criteria

- Babies with gestational age at birth of less than 32 weeks.
- Babies who were admitted to a neonatal unit within an hour of birth.
- Babies who experienced their final neonatal discharge in the calendar year of analysis.
- Only the first known episode of care will be considered for analysis.

Attribution

Results will be reported for each hospital of birth and network of birth. When the place of birth is unknown, a non-hospital location, or a hospital without a neonatal unit, the place of birth will be assigned as “Other”.

Deriving outcomes

NNAP will use data from the first temperature taken after admission to categorise when babies had their temperature taken, and the value of temperature measurement (Table 5).

Table 5: Categorising temperature measurement from BadgerNet fields *Temperature measured after admission, Temperature value* and *Temperature not recordable*

Time of first temperature measurement (minutes from birth)	Value of first temperature measurement				
	Temp. not taken	< 35.9	36-36.4	36.5-37.5	>37.5
Temperature not taken after admission	Not taken	Not taken	Not taken	Not taken	Not taken
Within an hour (≤ 60 minutes from birth)	Not taken	Within an hour	Within an hour	Within an hour and between 36.5-37.5	Within an hour and >37.5
After an hour (> 60 minutes from birth)	Not taken	Late	Late	Late	Late

Categorising temperature values

Babies with a temperature measured *Within an hour* will be further categorised into the following groups based on their temperature values:

- Less than 32.0°C
- 32.0°C-35.9°C
- 36.0°C-36.4°C
- 36.5°C-37.5°C (normothermic)
- Greater than 37.5 °C

When the temperature value was not recordable, babies will be assigned to the 32.0°C-35.9°C group.

BadgerNet data source

Details on temperature measurement are captured on BadgerNet on the *Details at Birth and Admission* page. Values from the *Temperature measured after admission, Temperature value* and *Temperature not recordable* fields are used by NNAP for analysis (Indicated below).




Figure 4: Details at birth and admission page, BadgerNet

Parental consultation within 24 hours of every admission

Is there a documented consultation with parents by a senior member of the neonatal team, within 24 hours, of admission.

11,12,17?

Note: By senior member of the neonatal team, NNAP means a consultant or middle grade doctor, or a nurse practitioner acting in such a role.

Change to audit measure for 2020 data year: This measure now considers every admission to a neonatal unit of 12 hours or longer for each baby, not just first admission as in previous years.

NNAP standard

Developmental standard: A consultation should take place within 24 hours of admission for every baby (100%).

Source of standard: NNAP Project Board

Comparison standard for outlier analysis: National rate.

Inclusion criteria

- Babies who experienced their final neonatal discharge in the calendar year of analysis.
- Babies who were admitted to neonatal care for at least 12 hours for a given episode, receiving special care or a higher order of neonatal care.
- Babies receiving neonatal care in non-neonatal unit locations (postnatal ward, transitional care etc.) will not be included.

Attribution

Results will be attributed to the neonatal unit and network of admission.

Deriving outcomes

The recorded time for first consultation with a senior member of staff is compared to the time of admission to determine when consultation takes place. The NNAP standard is aimed specifically at consultation at or after the time of admission, and so consultations prior to admission or birth cannot meet the NNAP standard (Table 6).

Table 6: Categorising consultation with parents from BadgerNet fields *Parents seen by senior staff* and *Time first seen*

Time of first consultation	NNAP category
Before admission	Before admission
Time of admission – 24 hours after admission	Within 24 hours (NNAP standard)
More than 24 hours after admission	After 24 hours
Confirmed “no consultation”	No consultation
Confirmed “unknown”	Missing / unknown data
Missing consultation time	Missing / unknown data

BadgerNet data source

Details on first consultation are captured on BadgerNet on the *Details at Birth and Admission* page. The answer to the question *Parents seen by senior staff* and the *Time first seen* are used by NNAP for analysis.




Figure 5: Details at birth and admission page, BadgerNet

Parental presence at consultant ward rounds

For a baby admitted for more than 24 hours, did at least one parent attend a consultant ward round^{11,12,13}?

Note: Consultant ward round refers to any ward round where a consultant is in attendance, at any time of the day.

Change to audit measure for 2020 data year: None

NNAP standard

Developmental standard: Benchmarking only.

Comparison standard for outlier analysis: Not applicable, no outlier analysis.

Inclusion criteria

- Babies who experienced their final neonatal discharge in the calendar year of analysis.
- Admissions will be included if there is at least 24 hours (≥ 1440 minutes) between the admission time and discharge time for the episode of care.
- It is possible for one baby to have multiple eligible admissions for this analysis.
- Babies receiving all neonatal care in non-neonatal unit locations (postnatal ward, transitional care etc.) will not be included.

Attribution

Attribution will be to the neonatal unit providing care in each eligible admission.

Deriving outcomes

NNAP will use data captured as part of the daily summary of care to confirm parental presence on consultant led ward rounds, and admissions will be classified based on aggregated attendance data (Table 7).

Table 7: Categorising parental presence at consultant ward rounds from BadgerNet field *Parent present on consultant ward round today?*

Number of days where a parent was present on consultant ward round	NNAP category
None	Parent(s) not present for any ward rounds during admission
One or more	Parent(s) present for one or more ward rounds during admission

Days where no data is entered concerning parental presence on ward rounds, or where daily data is missing, will be considered as parental absence from the ward round.

BadgerNet data source

Details on daily parental presence on ward rounds are captured on BadgerNet in daily summary forms in the *Parent present on consultant ward round today?* field. A form can be created for each baby for each calendar day where they are an inpatient, and NNAP will consider all of the forms created in relation to each eligible admission when determining parental presence.

The screenshot shows the 'General Details' section of a BadgerNet form. On the left is a sidebar with various medical categories. The main form area contains the following fields and options:

- Patient care date:** 01 Nov 16
- Location of care today:** Radio buttons for NNU, TransCare, PN ward, and Other obstetric area.
- Ward location:** A dropdown menu.
- Reduced form for TC or PNWARD data:** A button labeled 'Show whole input form'.
- Weight today:** An input field followed by 'grams' and the note '(complete only if baby actually weighed today)'. Below it is a 'Get latest weight' button.
- Working weight:** An input field followed by 'grams'.
- Most recent head circumference:** An input field followed by 'cm'.
- Most recent length:** An input field followed by 'cm'.
- Requiring 1:1 nursing today:** Radio buttons for 'Yes' and 'No (sick/unstable)'.
- Carer Status:** Radio buttons for 'Carer resident - Caring for baby', 'Carer resident - Not caring for baby', and 'Carer not resident'.
- Nursing Status:** Radio buttons for 'Barrier nursed' and 'Isolation'.
- Observations/Monitoring:** Radio buttons for 'None or >4hrly intervals', 'Continuous', 'Special', and 'Obs at regular intervals'.
- Parent present on consultant ward-round today?:** Radio buttons for 'Yes - Parent was present', 'No - Parent was not present', and 'No - no consultant ward-round'. A red arrow points to this field.
- Transported today:** Radio buttons for 'With Nurse Only', 'With Nurse and Doctor', 'With Paramedic', and 'Not transported'.

Figure 6: Daily summary, BadgerNet

Additional information

Background

The parents of babies admitted for care in neonatal units find themselves in a difficult and stressful situation, feeling that they have very little control in, or input towards, the care of their baby. It is therefore crucial that neonatal unit staff continue to keep parents informed of how their babies are being cared for and also listen to parents, try to understand how they are feeling and respond to any questions that they may have. This continual dialogue will also help parents to prepare for discharge, and transfer if required, by understanding the progress their babies have made and the progress they need to make prior to these stages.

This question is in addition to the consultation with parents within 24 hours of admission question.

Opportunities for quality improvement

The proposed metric will allow units to compare a measure of daily parental involvement with similar types of unit and might allow units to reflect on whether different levels of parental involvement in ward rounds might improve parental partnership in care.

On-time screening for retinopathy of prematurity

Does an admitted baby born weighing less than 1501 g, or at gestational age of less than 32 weeks, undergo the first retinopathy of prematurity (ROP) screening in accordance with the NNAP interpretation of the current guideline recommendations¹⁴?

Change to audit measure for 2020 data year: None.

NNAP standard

Developmental standard: All (100%) of eligible babies should receive ROP screening within the time windows for first screening recommended in the guidelines.

Source of standard: National standard (RCPCH, RCOphth, BAPM and Bliss, *Guideline for the Screening and Treatment of Retinopathy of Prematurity, 2008*¹⁴).

Comparison standard for outlier analysis: National rate

Note: In interpreting the national standards for this NNAP analysis, the Project Board has decided that a baby will be seen as having had ROP screening “on-time” if:

- *A baby who was discharged before the ROP screening window opened had their first screening conducted prior to discharge, or*
- *A ROP screen takes place within the ROP screening window, before or after discharge.*

The NNAP Project Board has also agreed to allow an extra week either side of the ROP screening window (Table 8).

Table 8: ROP screening windows

Gestational age at birth (completed weeks)	ROP screening windows	
	National guideline ROP screening window	NNAP ROP screening window
Less than 27	30 ⁺⁰ to 30 ⁺⁶ weeks corrected gestational age inclusive	29 to 31 weeks corrected gestational age inclusive
Greater than or equal to 27	4 to 5 weeks from birth (28-35 days)	3 to 6 weeks from birth (21-42 days)

Inclusion criteria

- Babies who experienced their final neonatal discharge in the calendar year of analysis
- The baby was alive at the beginning of the national guideline screening window

and

- The baby was born at less than 32 weeks gestational age and was admitted to a neonatal unit

or

- The baby's birth weight was 1500 g or less.

Attribution

If a ROP screen is conducted in accordance with the NNAP standard this is assigned to the neonatal unit performing the first screen within the NNAP ROP screening window.

If a baby is not screened in accordance with the NNAP standard this is assigned to the neonatal unit of care at the time that the national guideline screening window (not the extended NNAP ROP screening window) closed, or the neonatal unit of final discharge when the infant was discharged before the closure of the national guideline screening window.

Babies who died before the end of screening window, and who do not have a record of being "Screened on time" will be not be attributed to a unit.

Deriving outcomes

ROP status will be derived from the ad -hoc and daily summary of care sections of BadgerNet to confirm if a screening took place and whether it took place during the NNAP ROP screening window (Table 9).

Table 9: NNAP ROP screening done categories

These categories are totalled to give 'Any screen':			No screening data
Within NNAP ROP screening window	Before NNAP ROP screening window opened	After NNAP ROP screening window	No ad-hoc ROP form or ROP screen data on daily care summary
Screened on time	Screened early	Screened late	No screening data

BadgerNet data source

Details on ROP screenings are captured on BadgerNet via ad-hoc forms for ROP screening and from the daily summary form when *ROP screen today* is answered *Yes*. Where evidence of ROP screening is available from both sources for the same day ad-hoc screening data will be used preferentially.

The *Date and time* from the ad-hoc form is used to determine the time of screening, whilst daily summary is considered to have happened at the very beginning of the day (00:00).

The figure displays two screenshots of the BadgerNet interface. The top screenshot shows an ad-hoc form for ROP screening for the 'Right Eye'. It includes fields for 'Date and time' (11 Jan 16 at 10:19), 'Highest ROP Stage in any Zone' (radio buttons for No ROP, One, Two, Three, Four, Five, AP-ROP), 'Regression of ROP' (radio button for Yes), 'Clock hours' (radio buttons for 0-12), 'Zone of vascularisation' (radio buttons for No ROP, Zone 1, Zone 2, Zone 3), 'Plus disease' (radio buttons for None, Pre-plus, Plus disease), and a 'Notes' text area. A red arrow points to the 'Date and time' field. The bottom screenshot shows a daily summary form with a sidebar menu on the left containing categories like 'General information', 'Respiratory', 'Cardiovascular', 'Gastrointestinal', 'Neurology/NAS', 'Ophthalmology', 'Lines in situ/Sepsis', and 'Metabolic/ Jaundice'. The main content area includes 'ROP screen today' (radio buttons for Yes, No), 'ROP treatment today' (radio buttons for Yes, No), 'Daily comments' text area, and 'ROP notes for discharge (nothing noted)' with an 'Update discharge notes...' button. A red arrow points to the 'ROP screen today' field.

Figure 7: Ad-hoc form for ROP screening (top) and daily summary form (bottom), BadgerNet

Bloodstream infection

Does an admitted baby have one or more episodes of bloodstream infection, characterised by one or more positive blood cultures taken, after 72 hours of age?

Change to audit measure for 2020 data year: None. From 2018, there was a change to the inclusion criteria to so that the denominator includes only babies still present on the neonatal unit at 72 hours of age.

NNAP standard

Developmental standard: Benchmarking only.

Comparison standard for outlier analysis: Not applicable, no outlier analysis.

Inclusion criteria

- Babies who experienced their final neonatal discharge in the calendar year of analysis.
- Babies who were present on the neonatal unit at 72 hours of age.

Attribution

Blood cultures will be attributed to the neonatal unit of care when the culture was taken, or for units where there were two such units on a day of care, the unit completing the culture form. Babies will be counted as eligible once in each neonatal unit they were admitted to.

Deriving outcomes

Culture records will be considered as complete if the pathogen results for the culture are entered including 'None' or 'No growth'.

Fungal, and bacterial culture growths will be categorised as pure growths of a clearly pathogenic organism, mixed growths or organisms of uncertain significance (including skin commensals) according to a predefined list of clearly pathogenic known pathogens (see Appendix 1 for the list used for 2017 data).

Data pertaining to late onset bloodstream infection will be presented in gestational age sub groups, and subdivided by positive cultures for "known pathogens" and "skin commensal" organisms (organisms which may represent "false positive" blood cultures).

Data viewing will be further enhanced by presentation of an indication that all positive blood cultures have been entered, for centres where this is the case.

BadgerNet data source

Details for blood cultures and their results will be captured via ad-hoc forms for cultures on BadgerNet.

The figure displays two screenshots of the BadgerNet ad-hoc form for blood culture. The top screenshot shows the 'Culture Details' form, and the bottom screenshot shows the 'Blood Result' form.

Top Screenshot: Culture Details

Culture Details

Blood Result

Date and Time Sample(s) Taken: 13 Nov 18 at 10:25

Sample Type(s)

- Blood culture
- CSF Culture
- Urine
- Secretions
- Swab
- Other

Decision to treat: at

Taken from: Central line Peripheral line Closed culture

Paired sample: Yes No

Reason for Culture: [Dropdown]

Signs Present When Culture(s) Obtained: [Dropdown]

Prep used: 2% Chlorhexidine 5% Chlorhexidine

HeRo score (if done): [Text]

Time of 1st antibiotics: [Dropdown] at [Text]

Bottom Screenshot: Blood Result

Blood Result

Pathogens No growth

Pathogens: [Dropdown]

Sensitivity List for Isolate: [Dropdown]

Notes: [Text Area]

Figure 8: Ad-hoc form for blood culture taken (top), and ad-hoc form for blood culture results (bottom)

Central line associated bloodstream infection (Quality Improvement Surveillance Definition)

How many babies have a positive blood culture (any species) with a central line present, after the first 72 hours of life, per 1000 central line days?

Change to audit measure for 2020 data year: None. From 2018, there was a change to the inclusion criteria to so that the denominator includes only babies still present on the neonatal unit at 72 hours of age.

NNAP standard

Developmental standard: None, benchmarking only.

Comparison standard for outlier analysis: Not applicable, no outlier analysis.

Inclusion criteria

- Babies who experienced their final neonatal discharge in the calendar year of analysis.
- Babies who were present on the neonatal unit at 72 hours of age.
- All days where a central line (surgical venous line, umbilical venous catheter (UVC), umbilical artery catheter (UAC), peripherally inserted central catheter (PICC)) was present will be included in the number of line days when calculating proportions per 1000 line days.

Attribution

Positive growths will be attributed to the neonatal unit completing the culture form and babies will be counted as eligible once in each neonatal unit they were admitted to.

Deriving outcomes

To ensure that early onset infections are not counted in error cultures from the first 72 hours of life will initially be included in the analysis. A blood culture that presents any growth will be considered positive, and recurring growths that are less than 72 hours apart will be removed so that only the earliest known instance of the growth remains in the data. Once this comparison has been complete, any remaining cultures from the first 72 hours of life will be removed from the analysis.

The time the culture was taken will then be used to map to daily data records to determine if a central line was present on the calendar day the culture was taken. If a daily record is not available for the day, the culture was taken then it will be assumed that a central line was not present.

BadgerNet data source

Details for blood cultures and their results will be captured via ad-hoc forms for cultures on BadgerNet, whilst data on *Lines in situ today* from daily data summaries will be used to determine line days for the analysis.

The top screenshot shows the 'Culture Details' form. The 'Date and Time Sample(s) Taken' is 13 Nov 18 at 10:25. The 'Sample Type(s)' section is highlighted in green and contains a checked 'Blood culture' option. Other options include CSF Culture, Urine, Secretions, Swab, and Other. The 'Decision to treat' is 'at'. 'Taken from' includes Central line, Peripheral line, and Closed culture. 'Paired sample' is 'No'. 'Reason for Culture' and 'Signs Present When Culture(s) Obtained' are dropdown menus. 'Prep used' includes 2% Chlorhexidine and 5% Chlorhexidine. 'HeRo score (if done)' and 'Time of 1st antibiotics' are also present.

The bottom screenshot shows the 'Blood Result' form. The 'Pathogens' field contains 'No growth'. There are also fields for 'Pathogens', 'Sensitivity List for Isolate', and 'Notes'.

Figure 9: Ad-hoc form for blood culture taken (top), and ad-hoc form for blood culture results (bottom), BadgerNet

The screenshot shows the 'Lines in situ/Sepsis' section. The 'Lines in situ today' dropdown menu is highlighted in green and shows a list of line types: None, Peripheral arterial line, Umbilical arterial line, Umbilical venous line, Percutaneous central venous line (long line), Surgically inserted central venous line in situ, Peripheral venous line in situ, and Subcutaneous. The right side of the form contains management and screening summary information.

Figure 10: Daily summary form, BadgerNet

Bronchopulmonary dysplasia

Does an admitted baby born at less than 32 weeks develop bronchopulmonary dysplasia (BPD) or die?

Change to audit measure for 2020 data year: None.

NNAP standard

Developmental standard: None, benchmarking only.

Comparison standard for outlier analysis: Treatment effect of 0% (network level only).*

**Note: BPD in a network is assessed by comparing its rate with the rate of a comparable set of babies in the UK (or NNAP) as a whole. More information about this method is found in the NNAP statistical analysis plan.*

Inclusion criteria

- Babies with a gestational age at birth of less than 32 weeks.
- Babies who experienced their final neonatal discharge in the calendar year of analysis.

AND

- Babies who were still an inpatient in a neonatal unit at 36 weeks postmenstrual age, had been discharged alive from neonatal care at less than 36 weeks postmenstrual age, or had died before 36 weeks postmenstrual age.

Attribution

Attribution will be to the hospital of birth. Outcomes will also be reported by network of birth. When the place of birth is unknown, a non-hospital location, or a hospital without a neonatal unit, the place of birth will be assigned as *Other*.

Deriving outcomes

Definition of bronchopulmonary dysplasia

BPD will be defined by the level of respiratory support received at 36 weeks postmenstrual age (Table 10).

When a baby was discharged from neonatal care before reaching 36 weeks corrected gestational age, data from the time of discharge will be used instead.

Table 10: Categorising BPD or death in NNAP based on daily respiratory support data

Survival at 36 weeks CGA	Respiratory data from 36 weeks corrected gestational age / final discharge		
	Not receiving respiratory support	Receiving respiratory support*	Missing required respiratory data**
Died before 36 weeks CGA	Died	Died	Died
Survived to 36 weeks CGA	No BPD	BPD	BPD not determinable

**Respiratory Support' is defined as receiving any of the following at the specified time:

- Ventilation
- CPAP
- Non-invasive ventilation (e.g. BiPAP)
- Hi flow nasal cannula therapy (i.e. ≥ 2 lpm gas flow)
- Oxygen treatment

** In instances where respiratory data is not available for the specified day, data from the subsequent day of care (or penultimate day of care for babies discharged less than 36 weeks) will be sought instead. When respiratory data is also missing from this second source the baby will be classified as 'Missing required data' for that period.

Babies will be defined as having missing data if there is no data available to determine respiratory support.

BadgerNet data source

Details on respiratory support used to calculate levels of BPD are captured on BadgerNet in daily summary forms. A form is created for each baby for each calendar day where they are an inpatient on a neonatal unit, and the fields *Respiratory Support*, *Added oxygen*, *Mode of ventilation* and *Mode of non-invasive support* will be used to by the NNAP.

The image shows a screenshot of the BadgerNet daily summary form. On the left, there is a vertical navigation menu with categories: General information, Respiratory, Cardiovascular, Gastrointestinal, Neurology/NAS, and Ophthalmology. The 'Respiratory' section is highlighted. To the right of the menu, there are four main input fields, each with a red arrow pointing to it from the 'Respiratory' menu item:

- Respiratory support:** A dropdown menu with three options: 'No ventilation or CPAP' (highlighted in green), 'Ventilation via ET tube / tracheostomy', and 'Non invasive support (inc CPAP)'.
- Added oxygen:** A dropdown menu.
- Mode of ventilation:** A dropdown menu with two options: 'Conventional' and 'HFOV'.
- Mode of noninvasive support:** A dropdown menu.

Below these fields, there is a checkbox for 'Nasopharyngeal Airway in situ' with 'Yes' and 'No' options.

Figure 11: Daily summary form, BadgerNet

Necrotising enterocolitis

Does an admitted baby born at less than 32 weeks gestational age meet the NNAP surveillance definition for necrotising enterocolitis (NEC) on one or more occasion?

Change to audit measure for 2020 data year: None.

NNAP standard

Developmental standard: Benchmarking

Comparison standard for outlier analysis: Not applicable, no outlier analysis.

Background and evidence base

Inclusion criteria

- Babies who experienced their final neonatal discharge in the calendar year of analysis.
- Babies born at less than 32 weeks gestational age and survived to at least 48 hours after birth.

Attribution

- Babies will be attributed to their location of care at 48 hours of life, which is intended as a proxy measure of the intention to provide ongoing care for a baby in a given neonatal unit.
- When a baby is in transit between units at 48 hours the baby will be assigned to the transferring hospital. When multiple admission locations exist at 48 hours of life, the baby will be attributed to the earliest associated admission time.

Deriving outcomes

The analysis is based upon where the baby was resident at 48 hours of age. NEC may be diagnosed at surgery, post-mortem or based on the following clinical and radiographic signs.

At least one clinical feature from:

- a) Bilious gastric aspirate or emesis
- b) Abdominal distension
- c) Occult or gross blood in stool (no fissure)

And at least one radiographic feature from:

- a) Pneumatosis
- b) Hepato-biliary gas
- c) Pneumoperitoneum

Infants clinically diagnosed as NEC using the clinical and radiographic criteria who are found at surgery or post-mortem to have "Focal Intestinal Perforation" should not be recorded as having NEC.

Table 11: Categorisation of NEC diagnosis

Survival to discharge home	Was NEC Diagnosed during any admission?					
	NEC diagnosis based on surgery	NEC diagnosis based on post-mortem	NEC diagnosis based on presence of clinical signs		NEC not diagnosed	NEC diagnosis data missing
			At least 1 radiographic and 1 clinical feature	Radiographic or clinical features missing		
Survived to discharge home	NEC	NEC	NEC	Missing data (alive at discharge)	No NEC	Missing data (alive at discharge)
Died prior to discharge home	NEC	NEC	NEC	Missing data (died before discharge)	No NEC but died prior to discharge	Missing data (died before discharge)

BadgerNet data source

For babies born at less than 32 weeks, the field *Was NEC diagnosed during this admission?* will appear on the *Discharge details*. Data from this field, and it's follow on question (*Based on, Clinical feature(s), Radiographic Features*).

The screenshot shows the 'Discharge questions' section of the BadgerNet interface. It contains the following questions and options:

- Oxygen at Discharge: Yes No
- Aproxae/Cardiorespiratory/Saturation Monitoring at Discharge: Yes No
- Did baby have any ROP screening this episode?: Yes No
- Was NEC diagnosed during this admission?: Yes No

The 'Based on' section for the NEC question is highlighted, showing:

- Clinical signs
- Surgical confirmation
- Post-mortem examination

Below this, there are sections for 'Clinical feature(s)' and 'Radiographic feature(s)', each with a list of features and checkboxes:

- Clinical feature(s):**
 - Bilious gastric aspirate or emesis
 - Abdominal distension
 - Occult or gross blood in stool (no fissure)
- Radiographic feature(s):**
 - Pneumatosis intestinalis
 - Hepato-biliary gas
 - Pneumoperitoneum

Figure 12: Discharge details page, BadgerNet

Additional information

Background

Necrotising enterocolitis (NEC) is a devastating consequence of preterm delivery typically affecting around 5% of babies born at less than 32 weeks gestation, with known variation between centres. Mortality is high, exceeding 20%. Observational evidence suggests that survivors experience substantially higher rates of developmental impairment. Hospital stays in survivors are typically prolonged, and frequently require transfer with prolonged care, often in surgical centres. As well as being disadvantageous for babies and families, this confounds analyses of incidence where cases are attributed to the hospital providing the majority of the care days.

Opportunities for quality improvement

NEC is known to be influenced by antenatal factors, such as in utero growth. The evidence base for the widely-held view that postnatal feeding strategies influence necrotising enterocolitis is, as yet, inconclusive. Limited evidence supports the assertion that alterations in feeding increments can moderate the risk of NEC. However, meta-analyses based on older trials of donor breast milk suggest that using formula milk substantially increases the risk of NEC. Finally, meta-analysis of probiotic trials in very large numbers of preterm babies suggests that probiotic use moderates the risk of NEC.

Measuring rates of NEC will therefore afford neonatal units opportunities to compare their rates of NEC with those of other neonatal units, in a way unaffected by variations in interventions rates according to local surgical practice. Units with higher NEC rates will wish to address their adherence to strands from known quality improvement initiatives.

It is known that a significant number of babies with NEC die without surgery - describing the site where NEC is first diagnosed potentially represents a quality improvement opportunity.

Minimising inappropriate separation of mother and term baby

For a baby born at greater than or equal to 37 weeks gestational age, who did not have any surgery or a transfer during any admission, how many special care^a or normal care^b days were provided when oxygen was not administered?

a= Healthcare Resource Group (HRG) 3 or b= HRG 5.

Change to audit measure for 2020 data year: None.

NNAP standard

Developmental standard: None, benchmarking only.

Inclusion criteria

- Babies born greater than or equal to 37 weeks.
- Babies who received all their care in one unit.
- Babies who were admitted for at least 12 hours.
- Babies who did not have major surgery.
- Babies who were nursed on a neonatal unit on a day, or days.
- Babies receiving neonatal care in non-neonatal unit locations (postnatal ward, transitional care etc.) will not be included.

A second denominator of all live births within the gestational age range in the hospital (greater than or equal to 37 weeks) will be used when available.

Attribution

Attribution will be to the neonatal unit of admission.

Deriving outcomes

NNAP will use the pre-calculated care levels from BadgerNet to determine care days that are classified as HRG 3 or HRG 5, with *no added oxygen* and no other form of non-invasive respiratory support provided. In addition, only care days that occurred in a neonatal unit location will be counted.

Days of care will be defined as having missing data if there is no data available to determine respiratory support.

BadgerNet data source

Details from the *Any major surgery today* field from daily data summaries will be used to determine and exclude babies who received surgery from the analysis.

Furthermore, the pre-calculated care level from each daily data summary, and the respiratory support details for that day, will be used to categorise care days.

Figure 13: Daily summary nursing form - diagnosis, procedures and drugs (top), general information (middle) and respiratory (bottom), BadgerNet

Additional information

Background

Between 2011 and 2014, the number of term live births in England declined by 3.6% yet the number of term babies admitted to the neonatal units increased by 31%. This increase is seen across all categories of care but especially marked in the special care category where almost 10,000 more babies were admitted for this type care in 2015 compared to 2011. If more facilities were available to nurse baby and mother together when babies require monitoring or intervention that is not part of “normal care”, it is possible that many such admissions and unnecessary separation of babies from their mothers can be avoided.

Opportunities for quality improvement

Analysis of retrospective patient level data for term neonatal admissions between 2011 and 2014 commissioned by NHS England suggests that a substantial proportion of these admissions may be preventable with different care model. For example, 23% of admissions for respiratory problems had a length of stay of less than 48 hours and needed special care only, with 10% not needing oxygen at all. Most babies admitted for jaundice required phototherapy only and could be more appropriately managed in a transitional care setting. Similarly, amongst babies admitted for hypoglycaemia, 30% of admissions occurred before four hours of age - a period of physiological transition in glucose metabolism (with half of these within an hour of birth), 44% admitted directly from the delivery suite, and admission blood glucose concentration being above the operational threshold in a quarter of these admissions, suggesting little or no postnatal ward interventions. This was supported by the finding that nearly 75% of babies admitted for hypoglycaemia within an hour of birth did not need intravenous glucose infusion.

A recent quality improvement initiative showed a 73% reduction in admissions of later preterm and term babies for hypoglycaemia by adhering to the currently available guidance¹⁵. This, together with the retrospective data analysis suggests opportunities for wider quality improvement and reductions in unnecessary separation of mothers and their babies.

This metric will:

- Be inclusive - all neonatal units irrespective of their designation will be able to participate in the resulting quality improvement
- Address the largest group of babies admitted to any neonatal unit (term neonates)
- Forge a genuine partnership in quality improvement including not only neonatal, obstetric and midwifery healthcare professionals but also active involvement of the parents.

Minimising inappropriate separation of mother and late preterm baby

For a baby born at 34-36 weeks gestational age, who did not have any surgery or a transfer during any admission, how many special care^a or normal care^b days were provided when oxygen was not administered?

a= HRG 3 or b= HRG 5.

Change to audit measure for 2020 data year: None.

NNAP standard

Developmental standard: Benchmarking only.

Inclusion criteria

- Babies born between 34 and 36 weeks gestational age.
- Babies who received all their care in one unit.
- Babies who were admitted for at least 12 hours.
- Babies who did not have major surgery.
- Babies who were nursed on a neonatal unit on a day, or days.
- Babies receiving neonatal care in non-neonatal unit locations (postnatal ward, transitional care etc.) will not be included.

A second denominator of all live births within the gestational age range in the hospital (34-36 weeks) will be used when available.

Attribution

Attribution will be to the neonatal unit of admission.

Deriving outcomes

NNAP will use the pre-calculated care levels from BadgerNet to determine care days that are classified as HRG 3 or HRG 5 with *no added oxygen* and no other form of non-invasive respiratory support provided. In addition, only care days that occurred in a neonatal unit location will be counted.

Days of care will be defined as having missing data if there is no data available to determine respiratory support.

The rate of potentially avoidable mother and baby separation days will be presented per admission. Data for each unit will also describe the number of admitted babies considered.

BadgerNet data source

Details from the *Any major surgery today* field from daily data summaries will be used to determine and exclude babies who received surgery from the analysis.

Furthermore, the pre-calculated care level from each daily data summary, and the respiratory support details for that day, will be used to categorise care days.

The figure displays three screenshots of the BadgerNet daily summary form, illustrating the data sources used for analysis.

Top Screenshot: Diagnosis, Procedures, and Drugs
 This section contains several dropdown menus for 'Diagnoses', 'Ongoing management issues today', 'Operations / Procedures today', and 'Investigations'. A red arrow points to the 'Any major surgery today' field, which has radio buttons for 'Yes' and 'No'.

Middle Screenshot: General Details
 This section includes fields for 'Patient care date', 'Location of care today', 'Ward location', 'Weight today', 'Working weight', 'Most recent head circumference', 'Most recent length', 'Requiring 1:1 nursing today', and 'Cater Status'. A red arrow points from the 'Weight today' field to the 'Clinical Summary' on the right, which provides a summary of the patient's care, including dates, location, and management details.

Bottom Screenshot: Respiratory
 This section contains fields for 'Respiratory support', 'Added oxygen', 'Mode of ventilation', 'Mode of noninvasive support', and 'Nasopharyngeal Airway in situ'. Red arrows point to the 'Respiratory support' dropdown (showing options like 'No ventilation or CPAP', 'Ventilation via ET tube / tracheostomy', and 'Non-invasive support (inc CPAP)'), 'Added oxygen', 'Mode of ventilation', and 'Mode of noninvasive support'.

Figure 14: Daily summary form - diagnosis, procedures and drugs (top), general information (middle) and respiratory (bottom), BadgerNet

Additional information

Background

Nationally, 20% of admissions to neonatal units are of babies between 34 and 36 weeks gestation, weighing more than 1800 g at birth. Not all such babies are nursed in neonatal units – practice varies according to clinical preference and availability of appropriate facilities. There are clear parent and organisational drivers to encourage clinically appropriate use of transitional care – now described by Healthcare Resource Group (HRG) definitions. There is a strong indication from NHS England that payment structures will favour delivery of HRG 4 (special care parent present) which has been reconfigured according to a consensus of current clinical practice.

Opportunities for quality improvement

The metric will allow benchmarking of the proportion of care days per relevant admission to a neonatal unit that would be potentially suitable for HRG 4 care. NNAP will not suggest by this metric that all such days would be suitable for HRG 4 care, but the measure will facilitate comparison between services, and highlight a group of babies of whom a subpopulation might benefit from the development of appropriate services, such as transitional care, or other provision of HRG 4 care. The inclusion of babies receiving intravenous fluids within this definition is deliberate, reflecting that some babies receiving intravenous fluids could tolerate and benefit from enteral feeds.

Early breastmilk feeding

Does a baby born at less than 32 weeks gestational age receive any of their own mother's milk at day 14 of life?

Change to audit measure for 2020 data year: None.

NNAP standard

Developmental standard: Benchmarking only.

Inclusion criteria

- Babies born at less than 32 weeks gestational age who survive to their 14th day of life.
- Babies who experienced their final neonatal discharge in the calendar year of analysis.

Attribution

- Babies will be attributed to their location of care at 48 hours of life, which is intended as a proxy measure of the intention to provide ongoing care for a baby in a given neonatal unit.
- When a baby is in transit between units at 48 hours the baby will be assigned to the transferring hospital. When multiple admission locations exist at 48 hours of life, the baby will be attributed to the earliest associated admission time.

Deriving outcomes

Babies will be classified as meeting the NNAP standard if they are noted to have received any of the following types of enteral feed on their 14th day of life.

- Suckling at the breast
- Mother's fresh expressed breastmilk
- Mother's frozen expressed breastmilk

BadgerNet data source

The primary source for information on feeding at discharge is daily summary data on enteral feeds for the 14th day of neonatal care on the BadgerNet system. When enteral feeding data is missing for the 14th day, data from 13th or 15th day of care will be used instead (the most positive result will be used).

The image shows a screenshot of the 'Fluids and feeding' section in the BadgerNet system. On the left is a sidebar with various medical categories: General information, Respiratory, Cardiovascular, Gastrointestinal, Neurology/NAS, Ophthalmology, Ines in situ/Sepsis, Metabolic/ Jaundice, Haem/transfusions, Renal/Genitourinary, Skin, Fluids and Feeding (highlighted), and Other Problems. The main content area is titled 'Fluids and feeding' and contains several input fields and buttons. At the top right of this section is a button labeled 'Get feeding information from yesterday...'. Below it are radio button options for 'Parenteral nutrition today (TPN)' (Yes/No) and 'IV glucose and electrolyte solutions' (Yes/No). A red arrow points to the 'Enteral feeds today' dropdown menu. Below this are radio button options for 'Probiotics' (Yes/No) and 'Was baby tube fed for any part of day?' (Yes/No). There are also dropdown menus for 'Method of feeding', 'Additives', and 'Mouth care today?'. At the bottom of the form is a text area for 'Daily comments'.

Additional information

Breastmilk confers benefit for preterm babies, both through reducing adverse outcomes such as necrotising enterocolitis, and also by long term effects on neurodevelopment.^{1,2,3} In order for babies to benefit from both early risk modification (e.g. reduction in NEC) and long-term benefits, mothers of very preterm babies have to be successful in establishing expression of breastmilk, and to sustain this expression and intent to breastmilk feed over a long period. The existing measure of breastmilk feeding within NNAP (prevalence of any breastmilk feeding at discharge home) assesses establishment of expression and its continuation to such a point where a baby can be discharged breastmilk feeding. This new measure is designed to assess the success of initiation of breastmilk expression, in order to facilitate comparison between units, and quality improvement activities based on this.

Opportunities for quality improvement:

Given the widely acknowledged importance of breastmilk in promoting improved outcomes among very preterm infants, units with lower rates of breastmilk feeding at 14 days will wish to compare their practices to those with higher rates of breastmilk feeding. Measures addressing practical, professional, environmental and peer support for breastmilk expression may be of relevance to unit and network based neonatal quality improvement teams.

Breastmilk feeding at discharge home

*Does a baby born at less than 32 weeks gestational age receive any of their own mother's milk at discharge to home from a neonatal unit?*¹³

Change to audit measure for 2020 data year: None. From the 2019 data year, change of gestational age upper limit to babies born at less than 32 weeks gestational age and change of inclusion criteria so that babies transferred during their neonatal care are no longer excluded.

NNAP standard

Developmental standard: Eighty percent (80%) of babies born at less than 32 weeks gestational age should receive at least some of their mother's milk at discharge home from the neonatal unit.

Source of standard: By consensus, in consultation with BAPM.

Comparison standard for outlier analysis: Not applicable, no outlier analysis.

Inclusion criteria

- Babies born at less than 32 weeks gestational age.
- Babies who are discharged home alive.
- Babies who experienced their final neonatal discharge in the calendar year of analysis.

Attribution

Attribution will be to the neonatal unit of final discharge.

Deriving outcomes

Babies will be classified as meeting the NNAP standard if they are noted to have received any of the following types of enteral feed on their final day of care:

- Suckling at breast
- Mother's fresh expressed breast milk
- Mother's frozen expressed breast milk.

Data will be presented subdivided by rates of breastfeeding, and breastmilk feeding by bottle.

BadgerNet data source

The primary source for information on feeding at discharge is daily summary data on enteral feeds for the last day of neonatal care on the BadgerNet system. When enteral feeding data is missing for the final day, data from the penultimate day of care will be used instead.

General information

Respiratory

Cardiovascular

Gastrointestinal

Neurology/NAS

Ophthalmology

Ines in situ/Sepsis

Metabolic/ Jaundice

Bleem/transfusions

Renal/Genitourinary

Skin

Fluids and Feeding

Other Problems

Fluids and feeding

Get Feeding information from yesterday...

Parenteral nutrition today (TPN) Yes No

IV glucose and electrolyte solutions Yes No

Enteral feeds today

Probiotics Yes No

Was baby tube fed for any part of day? Yes No

Method of feeding

Additives

Skin to skin or Kangaroo care today? Yes No

Mouth care today?

Daily comments

Figure 15: Daily summary form, BadgerNet

Follow-up at two years

Does a baby born at less than 30 weeks gestational age receive medical follow-up at two years gestationally corrected age (18-30 gestationally corrected age range of acceptable ages)?

Does a baby have complete results of a structured assessment recorded?^{4,16}

Change to audit measure for 2020 data year: None.

NNAP standard

Developmental standard: Ninety percent (90%) of babies with two-year follow-up data entered.

Source of standard: NNAP Project Board

Comparison standard for outlier analysis: National rate.

Inclusion criteria

- Babies born at less than 30 weeks who are not recorded as deceased within their episodic data (including final neonatal outcome).
- The eligible cohort runs for births from July to June each year, for babies who would have reached two years corrected age since the last annual report. Current and projected denominators are shown in Table 12.

Table 12: Categorising eligible babies for follow-up at two years of age

NNAP data year	Two-year assessment cohort: Time of birth (inclusive)
2018	July 2015 to June 2016
2019	July 2016 to June 2017
2020	July 2017 to June 2018
2021	July 2018 to June 2019

Attribution

Attribution will be to the neonatal unit of final discharge. When the unit of final neonatal discharge home cannot be ascertained, two-year data is assigned to the last known episode of neonatal care.

Deriving outcomes

Assessment data will be considered for analysis if it took place between 18 and 30 months corrected age for the infant. This is calculated using assessment date, date of birth and gestational age at birth. If any of these fields are blank, the infant will be classified as *Outside of date range*. If the *Reason child not assessed* field is filled in, any data entered for this record will not count towards the analysis (Table 13).

Table 13: Categorising two-year follow-up data

Two-year follow-up outcome form	NNAP category
Any two-year follow-up health data entered	Health data entered
Lost to follow-up	No health data entered: Lost to follow-up
Died post discharge	Health data entered
Responsibility of another unit	No health data entered: Not assessed for other reason
Local decision not to follow-up	No health data entered: Not assessed for other reason.
Empty follow-up form / No follow up form	No health data entered: No health data entered at all
Assessment occurred outside 18-30 months corrected gestational age	No health data entered: No health data entered at all

Assessment data is categorised based on questions from the [TRPG/SEND/NNAP 2-Year Corrected Age Outcome Form](#). Questions are determined as being related to severe or mild/moderate impairment, and impairment for babies is classified based on the highest order of impairment they show within each NNAP impairment category.

Table 14: Questions from the outcome form are associated with the following NNAP impairment categories

NNAP impairment category	Domains used from TRPG/SEND/NNAP form	Questions related to mild/moderate impairment	Questions related to severe impairment
Neurodevelopmental	1.Neuromotor	a. Does the child have difficulty walking? b. Is the child's gait non-fluent or abnormal reducing mobility? d. Is this child unstable or needs to be supported when sitting? f. Does this child have any difficulty with the use of one hand? g. Does this child have difficulty with the use of both hands?	c. Is this child unable to walk without assistance? e. Is this child unable to sit? h. Is this child unable to use hands (i.e. to feed)?
	6.Neurology	a. Has this child had a fit or seizure in the past 12 months? b. Is this child on any anticonvulsants? d. Has this child ever had ventriculo-peritoneal shunt inserted?	c. Has this child had more than 1 seizures a month despite treatment
	8.Development	a. Is the child's development between 3-6 months behind corrected age? b. Is the child's development between 6-12 months behind corrected age?	c. Is the child's development more than 12 months behind corrected age?
	9.Neurossensory	a. Does this child have a hearing impairment? b. Does this child have hearing impairment corrected by aids? d. Does this child have any visual problems (including squint)? e. Does this child have visual defect that is not fully correctable?	c. Does this child have hearing impairment not correctable with aids? f. Is this child blind or sees light only?
	10.Communication	a. Does this child have any difficulty with communication? b. Does this child have difficulty with speech (< 10 words/signs)? d. Does this child have difficulty with understanding outside of familiar context?	c. Does this child have < 5 meaningful words, vocalisations or signs? e. Is this child unable to understand words or signs?
Respiratory	3. Respiratory & CVS system	a. Does this child have limited exercise tolerance with or without treatment?	b. Does child require supplemental oxygen or other respiratory support?
Gastro-intestinal	4. Gastro-intestinal Tract	a. Is this child on a special diet? b. Does this child have a stoma?	c. Is this child having renal dialysis or awaiting renal transplant?

Categorising missing assessment data

Missing data for individual assessment questions can lead to an infant's developmental outcome being classified as *not determinable*. This occurs when the missing data still has the potential to change the impairment status of the infant. For instance, if an infant is classified as having *mild/moderate* impairment from their entered data there will be no consequence from missing data for other questions associated with mild / moderate impairment. If, however, the same infant was missing an answer to a severe impairment question the status of the infant would be *not determinable*, as the missing data could still impact their impairment classification. Therefore, an infant must have complete data within an NNAP category in order to be classified as having *no impairment*.

Provision of standardised health assessments

An additional analysis is conducted to determine how many babies with health data entered at 18-30 months corrected age were provided with a Schedule of growing, Bayley III or Griffiths assessment. Any data entered in relation to these standardised assessments will be considered as evidence of a standardised assessment being provided.

BadgerNet data source

Data from the two-year follow-up form will be used to analyse assessment data. Which sections are used for each part of the analysis is indicated below by the coloured arrows:

-  Assessment data and neurodevelopmental outcomes
-  Respiratory outcomes
-  Gastro-intestinal outcomes
-  Standardised assessment data

Figure 16: Two-year follow-up form, BadgerNet

Mortality to discharge in very preterm babies

Does a baby born at less than 32 weeks gestational age die before discharge home, or 44 weeks post-menstrual age (whichever occurs sooner)?

Note: Babies discharged to a hospice for palliative care (such as for compassionate extubation) and who die prior to 44 weeks post-menstrual age are counted in the numerator.

Change to audit measure for 2020 data year: None

NNAP standard

Developmental standard: None

Comparison standard for outlier analysis: To be confirmed.

Inclusion criteria

Denominator: Number of babies admitted to a neonatal unit whose birth gestation was 24 to 31 weeks gestation inclusive.

Numerator: Deaths of babies 24 to 31 weeks gestation inclusive, before discharge from hospital to home, or discharge for palliative care with agreed non-intervention plan in place followed by death prior to 44 weeks post-menstrual age (e.g. ventilated baby discharged to a hospice). In hospital deaths in units not submitting data to the NNAP will be included. Mortality at or after 44 weeks gestationally corrected age will be excluded.

Table 15: Categorising eligible babies for Mortality until discharge or 44 weeks PMA

NNAP data year	Cohort: Date of birth
2018	1 July 2015 to 30 June 2018
2019	1 July 2016 to 30 June 2019
2020	1 July 2017 to 30 June 2020
2021	1 July 2018 to 30 June 2021

Attribution

The NNAP will report mortality on three year rolling epochs. Attribution will be to network of birth.


Deriving outcomes

Raw and adjusted mortality rates will be published for network and national data, but not for units.


Case mix adjustment: For limited variables, including gestation, gender, multiplicity and ethnicity but neither antenatal steroid administration nor congenital anomalies.

BadgerNet data source

Use the *Final Neonatal Outcome* page to record deaths that occur at Non-NNAP units, along with the date and time of death. Please ensure that you click save and close to ensure that any outcomes recorded by another unit using BadgerNet but not participating in the NNAP are recorded.



The NNAP mortality analysis will only include babies admitted to neonatal units, in the first instance. However, we would like units to report liveborn but not admitted deaths using Badger at gestations down to, and including, 23 weeks. Only the briefest of details are required to complete a “delivery room death” admission in place of a “neonatal unit admission”.



Additional information

Rationale

The National Advisory Group on Clinical Audits and Enquiries (NAGCAE) suggested NNAP report mortality in 2013. The NNAP agreed to form an expert group, and has subsequently

discussed reporting with the MBRRACE-UK group. MBRRACE-UK report neonatal mortality (<28 days) on all liveborn babies including those not admitted for neonatal care and do not have plans to report later mortality at this point. NNAP mortality reporting will build on the firm foundations established by NDAU mortality reporting to date.

Mortality is a vitally important outcome of neonatal care: reporting will add to the NNAP's perspective on neonatal services. In the first phase the NNAP plans to report on mortality to discharge of admitted babies. This definition may facilitate quality improvement of care within the neonatal unit. One reason for omitting babies born at 23 weeks gestation from the initial phase of reporting is that there are known important variations in rates of admission of babies at very low gestations. Low gestation babies have the highest mortality. Starting by reporting only on babies of 24-31 weeks gestation inclusive will maximise the comparability of reported neonatal network mortality to discharge.

Future phases of mortality reporting

Once NNAP mortality reporting is established, it is anticipated that future phases of reporting will include numerators and denominators extending to 23 weeks gestation, and the inclusion of liveborn, but not admitted cases of mortality.

What next?

Neonatal services should have systems for ensuring reporting to NNAP, via Badger, of any pre-44 week gestationally corrected age mortality to babies born at 24 to 31 weeks gestation who were discharged to non NNAP reporting units (surgical units, hospices, paediatric wards).

Neonatal should have systems to report to NNAP, via Badger, liveborn but not admitted babies at 23-31 weeks gestation (inclusive) who later die. These will be small in number, but are significant to research projects, MBRRACE-UK and future NNAP mortality reporting.

Nurse staffing on neonatal units

- *What proportion of nursing shifts are numerically staffed according to guidelines and service specification?*
- *What proportion of shifts have sufficient staff qualified in speciality (QIS)?*
- *How many additional nursing shifts are required to be worked to meet guidelines and service specification?*

Note: Shifts are based on corresponding level of cot occupancy calculated by Badger system.

Measure one - Proportion of shifts numerically staffed according to guidelines and service specification

Numerator: Number of shifts where nurse staffing met or exceeded service specification rules (1:1 intensive care; 1:2 high dependency care; 1:4 special care; additional shift coordinator)^{4,17,18}.

Denominator: Number of shifts (data collection instrument and measure based on a two-shift model of each calendar day).

Measure two - Proportion of shifts with sufficient staff qualified in speciality (QIS)

Numerator: Number of shifts in which at least 70% of registered staff on duty were qualified in speciality.

Denominator: Number of shifts (data collection instrument and measure based on a two-shift model of each calendar day).

Measure three - Number of additional nursing shifts required to meet guidelines and service specification

This number of shifts is calculated by subtracting the number of nurses that were working on each shift, from the number of nurses that are required under the terms of the service specification, and summing the rounded total over the reporting period^{4, 17,18}.

Change to audit measure for 2019 data year: New measure for 2018 data year.

NNAP standard

Developmental standard: 100% of shifts staffed according to guidelines and service specification.

Inclusion criteria

A “nurse” is a member of non-medical staff with an allocated patient workload, or acting as a coordinator, with or without Nursing and Midwifery Council (NMC) registration, with or without a qualification in speciality. Staff members without allocated patients, such as housekeepers, research nurses, nurses in primarily educational roles are specifically excluded.

If the supernumerary shift coordinator is QIS, they are included in the count of nurses QIS in measure 2.

When entering data about partially worked day shifts, units should enter early or late shifts as follows: where a nurse is only present for an “early” or “late” part of a day shift, record as 0.5 shifts.

Any shifts with no cot occupancy or nurse numbers data will be removed from the measure.

Additionally, any units with less than or equal to 25% data completeness across all their entered shifts will be removed from the measure. This completeness is assessed based on the following items:

- Registered Nurses, qualified in specialty
- Registered Nurses, currently undertaking QIS
- Registered Nurses not qualified in specialty
- Non registered Nurses
- Supernumerary shift leader

Units where more than 50% of shifts are staffed with three registered nurses or fewer are excluded from measure two.

Use the guidance [Safe, sustainable and productive staffing: An improvement resource for neonatal care](#) to support data completion.

Attribution

Attribution will be to the neonatal unit.

BadgerNet data source

Data from the Nursing numbers update form (Figure 17) in BadgerNet will be used for this measure. The Nursing numbers update form is found within the Unit report section.

Neonatal Unit Nursing Numbers

Nursing numbers update

Nursing numbers for 01 Nov 17 night period
For nursing period 01 Nov 17 at 20:00 to 02 Nov 17 at 08:00.

Involved in direct patient care

Registered nurses, qualification in speciality: 8.0

Registered nurses, currently undertaking QIS training: 1.0

Registered nurses, not qualified in speciality: 1.0

Non registered nurses: 1.0

Supernumerary shift team leader: Yes No

Total: 12.0

Of the total how many were bank staff?: 3.00

Supernumerary to patient care

Other nurses on duty but not involved with care: 0.0

Number of ANNPs on duty: 0.0

Comments

Min: 7 0 4 1 1 12 0 6 10 12 0 0 15 -3(0)

Figure 17: Nursing numbers update form, BadgerNet

Data entered into the form is summarised in the Neonatal Unit Nursing Numbers report (Figure 18).

Period	Nurses caring for patients					Cot occupancy			Nurses required				QIS Required (IC & HD)				
	With qual.	Undertaking QIS	Without qual.	Not Reg.	Team Leader	Total	Bank Staff	IC	HD	SC	Admissions	Discharges	RAPM Recommends	Difference	Toolkit recommends	Difference	
01 Nov 17	Day	6	0	4	1	1	12	1	3.3	7	11.4	2	2	10.6	1.4(0) (13%)	6.8	-0.8
	Night	8	1	1	1	1	12	3	4	7	10.3	0	1	11.1	0.9(0) (8%)	7.5	1.5
02 Nov 17	Day	5	0	5	2	1	13	0	4	8.2	10.3	5	0	11.7	1.3(0) (11%)	8.2	-3.2
	Night	8	1	2	0	1	12	3	5.8	8	12.3	2	0	13.9	-1.9(0) (-13%)	9.8	-0.8
03 Nov 17	Day	5	0	7	0	1	13	0	5	8.3	10.4	2	2	12.7	0.3(0) (2%)	9.1	-4.1

Figure 18: Neonatal unit nursing numbers, BadgerNet

Additional information

Background

Neonatal units in England are commissioned by NHS England specialist commissioning, according to the service specification^{4,17,18}. Services in the devolved nations are commissioned on a comparable basis according to the related British Association of

Perinatal Medicine (BAPM) standards¹⁸. Nurse staffing is known to be associated with outcome, with higher levels of nurse staffing associated with improved outcomes¹⁹. Audit users have asked that BAPM report on measures of nurse staffing, and the NNAP Project Board agreed to report a comparative measure of nurse staffing to allow neonatal units and neonatal networks to identify opportunities to improve the planning and delivery of neonatal unit care.

Opportunities for quality improvement

It is known that neonatal unit staffing varies, and that some units experience significant challenges in delivering a staffing model that meets with the standards identified in the service specification⁴. This is in part due to the unplanned nature of neonatal care, and random variations in demand. Additionally, challenges in appointing, retaining and providing career progression to a very highly specialist workforce may result in the provision of fewer nurses on individual nursing shifts than are mandated by the service specification. Significant resources are available to help services plan their workforces to meet with the standards.

The measure describes the proportion of nursing shifts which are staffed in accordance with the service specification, and additionally for each period will report how many additional nurse shifts would have been required to provide staffing at least adhering to the service specification, assuming a completely flexible additional workforce.

The NNAP recognises that it is not likely that reporting data will result in immediate moves to fully adherent staffing of all shifts in all units. Rather it is intended that this measure will enable commissioners and providers to focus on efforts to improve staffing in units with the highest deficits.

NNAP glossary and abbreviations

ATAIN	Avoiding Term Admissions into Neonatal units
BAPM	British Association of Perinatal Medicine
BPD	Bronchopulmonary dysplasia
COP	Clinical Outcomes Publication
CSF	Cerebrospinal fluid
EPR	Electronic patient record
HQIP	Healthcare Quality Improvement Partnership
HRG	Healthcare resource group
Hyperthermia	A body temperature more than 37.5°C
Hypothermia	A body temperature less than 36.5°C
LNU	Local neonatal unit
MBRRACE-UK	Mothers and Babies: Reducing Risk through Audits and Confidential Enquiries across the UK
MCN	Managed Clinical Network
NCAB	National Clinical Audit Benchmarking
NCAPOP	National Clinical Audit and Patient Outcomes Programme
NDAU	Neonatal Data Analysis Unit
NEC	Necrotising enterocolitis
NICU	Neonatal intensive care unit
NMC	Nursing and Midwifery Council
NMPA	National Maternity and Perinatal Audit
NNAP	National Neonatal Audit Programme
NNRD	National Neonatal Research Database
NUU	Neonatal unit
Normothermia	A body temperature between 36.5°C and 37.5°C
ODN	Operational delivery network
PICC	Peripherally inserted central catheter
RCM	Royal College of Midwives
RCOG	Royal College of Obstetrics and Gynaecology
RCPCH	Royal College of Paediatrics and Child Health
RCOphth	Royal College of Ophthalmologists
ROP	Retinopathy of prematurity
SCU	Special care unit
UAC	Umbilical artery catheter
UVC	Umbilical venous catheter

Appendix 1: “Pathogens” in the NNAP

Bacterial, fungal and yeast positive blood cultures reported to the NNAP for the Bloodstream infection and Central line associated bloodstream infection measures have been classified as shown below into organisms whose growth would be regarded as indicative of a bloodstream infection without further confirmatory evidence, and into a list of other organisms. This list of organisms included for NNAP reporting is available below. We are grateful to Professor Paul Heath* for reviewing and updating this list.

**Professor of Paediatric Infectious Diseases and Honorary Consultant, Paediatric Infectious Diseases Research Group; Director, St Georges Vaccine Institute.*

Table 1: Bacterial fungal and yeast organisms included in NNAP reporting.

Clearly pathogenic organisms		
Achromobacter sp	Enterobacter sp.	Pseudomonas sp.
Acinetobacter baumannii	Enterococcus faecalis	Pseudomonas stutzeri
Acinetobacter lwoffii	Enterococcus faecium	Roseomonas mucosa
Acinetobacter sp.	Enterococcus sp.	Rothia spp
B haemolytic streptococci	Escherichia coli	S. Aureus
Bacillus cereus	Flavimonas oryzihabitans	Salmonella sp.
Beta-haemolytic strep. Group b	Flavobacterium sp.	Salmonella unnamed
Burkholderia capeczia	Gbs	Serratia
Burkholderia sp.	Gemella haemolysans	Serratia liquefaciens
Candida albicans	Gemella morbilarum	Serratia marcescens
Candida dubliniensis	Group b streptococcus	Serratia sp.
Candida glabrata	Haemophilus influenzae	Staphylococcus aureus
Candida guilliermondii	Haemophilus parainfluenzae	Staphylococcus capitis
Candida kefyr	Klebsiella aerogenes	Staphylococcus epidermidis
Candida krusei	Klebsiella oxytoca	Staphylococcus haemolyticus
Candida parapsilosis	Klebsiella pneumoniae	Staphylococcus hominis
Candida sp.	Klebsiella sp.	Staphylococcus warneri
Candida tropicalis	Lactobacillus sp.	Stenotrophomonas maltophilia
Citrobacter freundii	Listeria	Streptococcus - group b
Citrobacter koseri	Listeria monocytogenes	Streptococcus agalactiae
Citrobacter sp.	Listeria sp.	Streptococcus bovis
Clostridium perfringens	Morganella morganii	Streptococcus milleri
Coliform	Mrsa	Streptococcus mitis

Corynebacterium diphtheriae	Neisseria meningitidis	Streptococcus pneumoniae
Eikenella corrodens	Pasteurella multocida	Streptococcus pyogenes
Enterobacter aerogenes	Pasteurella sp.	Streptococcus viridans
Enterobacter agglomerans	Proteus mirabilis	Treponema pallidum
Enterobacter cloacae	Pseudomonas aeruginosa	Yeast

Other organisms		
Abiotrophia adiacens	Corynebacterium ulcerans	Ralstonia sp.
Acanthamoeba sp.	Diphtheroids	Roseomonas gilardii
Acid fast bacilli	Gram positive bacilli	Scopulaiopsis brevicaulis
Actinomyces bovis	Gram positive cocci	Shigella sonnei
Actinomyces sp.	Group g streptococcus	Sphingomonas
Aerococcus sp	Haemophilus sp.	Staph saprophyticus
Aerococcus viridans	Kocuria species	Staphylococcus - coagulase negative
Alcaligenes faecalis	Lactococcus sp.	Staphylococcus - coagulase negative (mixed)
Alpha haemolytic streptococci	Micrococcus luteus	Staphylococcus simulans
Anaerobes	Micrococcus sp.	Staphylococcus sp.
Bacillus sp.	Mixed growth	Stomatococcus mucilaginosus
Bacteroides sp.	Moraxella catarrhalis	Streptococcus - group g
Campylobacter sp.	Moraxella sp.	Streptococcus anginosus
Chryseobacterium sp.	Mycobacterium sp.	Streptococcus oralis
Clostridium sp.	Mycoplasma hominis	Streptococcus parasinguinis
Coagulase negative staphylococcus	Neisseria sp.	Streptococcus salivarius
Cons	Nocardia asteroides	Streptococcus sanguis
Cons (mixed)	Peptostreptococcus sp.	Streptococcus sp.
Corynebacter	Prevotella sp.	Toxoplasma gondii
Corynebacterium bacilli	Propionibacterium sp	Ureaplasma
Corynebacterium sp.	Propionibacterium acnes	Ureaplasma sp.
Corynebacterium striatum	Psychrobacter phenylpyruvicus	Yeasts (other)

References

- ¹ The AHSN Network. *PReCePT*. Available at: <http://www.ahsnnetwork.com/about-academic-health-science-networks/national-programmes-priorities/precept/>
- ² Oddie S, Tuffnell D J, McGuire W. Antenatal magnesium sulfate: Neuro-protection for preterm infants. *Archives of Disease in Childhood - Fetal and Neonatal Edition* 2015;100:F553-F557.
- ³ Royal College of Obstetricians and Gynaecologists. Magnesium Sulphate to Prevent Cerebral Palsy following Preterm Birth (Scientific Impact Paper No. 29). 2011. Available at: <https://www.rcog.org.uk/en/guidelines-research-services/guidelines/sip29/>.
- ⁴ NHS England. Neonatal Critical Care Service Specification. 2016. Available at: <https://www.england.nhs.uk/commissioning/wp-content/uploads/sites/12/2015/01/e08-serv-spec-neonatalcritical.pdf>.
- ⁵ World Health Organization. WHO recommendations for the prevention and treatment of postpartum haemorrhage. Geneva: WHO, 2012. Available at: https://www.who.int/reproductivehealth/publications/maternal_perinatal_health/9789241548502/en/
- ⁶ McDonald SJ, Middleton P, Dowswell T, Morris PS. Effect of timing of umbilical cord clamping of term infants on maternal and neonatal outcomes. *Cochrane Database of Systematic Reviews* 2013, Issue 7. Art. No.: CD004074. DOI: 10.1002/14651858.CD004074.pub3. Available at: https://www.cochrane.org/CD004074/PREG_effect-timing-umbilical-cord-clamping-term-infants-mother-and-baby-outcomes
- ⁷ Fogarty M, Osborn DA, Askie L, Seidler AL et al. Delayed vs early umbilical cord clamping for preterm infants: a systematic review and metaanalysis. *Am J Obs and Gyne* 2018;218:1-18. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29097178>
- ⁸ Tarnow-Mordi W, Morris J, Kirby A, Robledo K, Askie L. Delayed versus Immediate Cord Clamping in Preterm Infants. *N Engl J Med*. 2017;377:2445-2455. Available at: <https://www.nejm.org/doi/full/10.1056/NEJMoa1711281>
- ⁹ Wyllie J, Perlman JM, Kattwinkel J, et al. Part 7: Neonatal resuscitation: 2015 International consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation* 2015;95:e169-201. Available at: https://pediatrics.aappublications.org/content/136/Supplement_2/S120.long
- ¹⁰ International Liaison Committee on Resuscitation. Temperature Maintenance in the Delivery Room. Available at: <https://volunteer.heart.org/apps/pico/Pages/PublicComment.aspx?q=589>
- ¹¹ Scottish Government. *Neonatal Care in Scotland: A Quality Framework*. 2013. Available at: <http://www.gov.scot/Resource/0041/00415230.pdf>.
- ¹² Welsh Health Specialised Services Committee, NHS Wales. *All Wales Neonatal Standards - 2nd Edition*. 2013. Available at: <http://www.wales.nhs.uk/document/219405>.
- ¹³ Bliss. Bliss Family Friendly Accreditation Scheme. 2015. Available at: <https://www.bliss.org.uk/Shop/bliss-family-friendly-accreditation-scheme-booklet>.
- ¹⁴ Royal College of Paediatrics and Child Health, Royal College of Ophthalmologists, British Association of Perinatal Medicine, BLISS. Guideline for the Screening and Treatment of Retinopathy of Prematurity. 2008. Available at: <https://www.rcophth.ac.uk/wp-content/uploads/2014/12/2008-SCI-021-Guidelines-Retinopathy-of-Prematurity.pdf>.

¹⁵ Stewart CE, Sage ELM, Reynolds P Supporting 'Baby Friendly': a quality improvement initiative for the management of transitional neonatal hypoglycaemia. *Archives of Disease in Childhood - Fetal and Neonatal Edition*. 2016;101:F344-F347.

¹⁶ National Institute for Health and Care Excellence. Developmental follow-up of children and young people born preterm. NICE guideline (NG72). 2017. Available at: <https://www.nice.org.uk/guidance/ng72>.

¹⁷ Department of Health. Toolkit for high quality neonatal services. 2009. Available at: http://webarchive.nationalarchives.gov.uk/20130123200735/http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_107845.

¹⁸ British Association for Perinatal Medicine. Service Standards for Hospitals Providing Neonatal Care (3rd edition). Available at: <https://www.bapm.org/resources/service-standards-hospitals-providing-neonatal-care-3rd-edition-2010>.

¹⁹ Watson S, Arulampalam W, Petrou S, Marlow N, Morgan A, Draper E, Modi N, On behalf of the Neonatal Data Analysis Unit (NDAU) and the Neonatal Economic, Staffing, and Clinical Outcomes Project (NESCOP) Group. The effects of a one-to-one nurse-to-patient ratio on the mortality rate in neonatal intensive care: a retrospective, longitudinal, population-based study. *Archives of Disease in Childhood - Fetal and Neonatal Edition*. 2016;101:F195-F200.

RCPCH Audits

Version 1.0, Published by RCPCH November 2019.

The Royal College of Paediatrics and Child Health is a registered charity in England and Wales (1057744) and in Scotland (SCO38299)